

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number
WO 01/36365 A2

(51) International Patent Classification⁷: C07C 59/64,
59/72, 59/90, 205/37, 311/53, 255/54, C07D 213/30,
C07C 215/14, C07D 215/14, 333/56, 215/18, 261/08,
277/24, 309/06, A61P 9/06, 5/14, A61K 31/192, 31/277

Novum, S-141 57 Huddinge (SE). **GORDON, Sandra**
[UY/SE]; Karo Bio AB, Novum, S-141 57 Huddinge (SE).

(21) International Application Number: PCT/EP00/11554

(74) Agents: **BANNERMAN, David, Gardner et al.**; With-
ers & Rogers, Goldings House, 2 Hays Lane, London SE1
2HW (GB).

(22) International Filing Date:
16 November 2000 (16.11.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9927056.3 17 November 1999 (17.11.1999) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **KARO
BIO AB** [SE/SE]; Novum, S-141 57 Huddinge (SE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MALM, Johan**
[SE/SE]; Korsangarvagen 53, S-142 40 Skogas (SE). **LIT-
TEN, Chris** [NZ/SE]; Stubbestigen 6, S-147 52 Tumba
(SE). **APELQVIST, Theresa** [SE/SE]; Sagstuvagen 2E,
3tr, S-141 49 Huddinge (SE). **HEDFORS, Asa** [SE/SE];
Domarvagen 3nb, S-141 62 Huddinge (SE). **BRANDT,
Peter** [SE/SE]; Karo Bio AB, Novum, S-141 57 Huddinge
(SE). **EDVINSSON, Karin** [SE/SE]; Karo Bio AB,

Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 01/36365 A2

(54) Title: THYROID RECEPTOR ANTAGONISTS FOR THE TREATMENT OF CARDIAC AND METABOLIC DISORDERS

(57) Abstract: This invention relates to novel compounds which are thyroid receptor ligands, preferably antagonists, and to methods for using such compounds in the treatment of cardiac arrhythmias, thyrotoxicosis and subclinical hyperthyroidism.

BEST AVAILABLE COPY

Thyroid Receptor Antagonists for the Treatment of Cardiac and metabolic disorders

FIELD OF THE INVENTION

This invention relates to novel compounds which are thyroid receptor ligands, preferably antagonists, and to methods for using such compounds in the treatment of cardiac arrhythmias, thyrotoxicosis and subclinical hyperthyroidism.

BACKGROUND OF THE INVENTION

Nuclear hormone receptors comprise a class of intracellular, mostly ligand-regulated transcription factors, which include receptors for thyroid hormones. Thyroid hormones exert profound effects on growth, development and homeostasis in mammals. They regulate important genes in intestinal, skeletal and cardiac muscles, liver and the central nervous system, and influence the overall metabolic rate, cholesterol and triglyceride levels, heart rate, and affect mood and overall sense of well being.

There are two major subtypes of the thyroid hormone receptor, TR α and TR β , expressed from two different genes. Differential RNA processing results in the formation of at least two isoforms from each gene. The TR α_1 , TR β_1 and TR β_2 isoforms bind thyroid hormone and act as ligand-regulated transcription factors. The TR α_2 isoform is prevalent in the pituitary and other parts of the central nervous system, does not bind thyroid hormones, and acts in many contexts as a transcriptional repressor. In adults, the TR β_1 isoform is the most prevalent form in most tissues, especially in the liver and muscle. The TR α_1 isoform is also widely distributed, although its levels are generally lower than those of the TR β_1 isoform. A growing body of data suggests that many or most effects of thyroid hormones on the heart, and in particular on the heart rate and rhythm, are mediated through the TR α_1 isoform, whereas most actions of the hormones on the liver, muscle and other tissues are mediated more through the β -forms of the receptor. It is believed that the α -isoform of the receptor is the major drive to heart rate for the following reasons: (i) tachycardia is very common in the syndrome of generalized resistance to thyroid hormone in which there are defective TR β -isoforms, and consequently high circulating levels of T₄ and T₃; (ii) tachycardia was observed in the only

described patient with a double deletion of the TR β gene (Takeda et al, J. Clin. Endocrinol. & Metab. 1992, Vol. 74, p. 49); (iii) a double knockout TR α gene (but not β -gene) in mice showed bradycardia and lengthening of action potential compared to control mice; (iv) western blot analysis of human myocardial TRs show presence of the TR α_1 , TR α_2 and TR β_2 proteins, but not TR β_1 .

If the indications above are correct, an α -selective thyroid hormone receptor antagonist that interacts selectively with the heart would offer an attractive alternative treatment of heart related disorders, such as atrial and ventricular arrhythmias.

Atrial fibrillation (AF) is the most common type of sustained arrhythmia encountered in primary care practice and is significantly more common in elderly patients, thus reflecting a reduction in the threshold for AF with age. Pharmacological treatment of AF involves the following types of anti-arrhythmic drugs according to Vaughan-Williams classification: (i) of class I such as disopyramide and flecainide (sodium channel blocker); (ii) of class III such as amiodarone (potassium channel blocker, prolongation of repolarization); (iii) of class IV such as verapamil and diltiazem (calcium channel blocker). Many patients are also subjected to electric cardioversions in order to convert atrial fibrillation into sinus rhythm. It should be noted that current therapies are associated with pro-arrhythmic risks and anti-arrhythmic agents often have insufficient efficacy partly because effective doses are limited by side-effects.

Ventricular arrhythmia, especially sustained ventricular tachycardia (VT) and ventricular fibrillation (VF) is the main cause of death associated with heart attack. Historically, three types of antiarrhythmic agents, class I agents, β -adrenergic blockers (class II), amiodarone and sotalol, appeared to offer the best scope for mortality reduction in patients with cardiac disease by preventing the occurrence of VT/VF.

The outcome of CAST (Cardiac Arrhythmia Suppression Trial, *N. Engl. J. Med.*, 321 (1989) 406-412) and its successor SWORD (Survival With Oral D-sotalol trial, 1994) created much concern regarding the potential of class I agents and sotalol. It was found that class I agents did not decrease mortalities in patient groups at risk for sudden cardiac death. For some

subsets of patients, class I agents even proved to increase mortality. The SWORD trial was stopped when sotalol proved to give higher death rate in patients, compared with the placebo. A consequence of these results is that the use of implantable defibrillators and surgical ablation have increased and that the trend in the industry has been towards the development of highly specific class III agents. Some of these channel blockers have been withdrawn from clinical development due to proarrhythmic effects and the subject remains under intensive debate. In this context it should be noted that amiodarone, despite its complex pharmacokinetics, mode of action (amiodarone is not regarded as a pure class III agent) and numerous side effects, is currently considered by many to be the most effective agent in the control of both atrial and ventricular arrhythmia.

Thyrotoxicosis is the clinical syndrome that results when tissues are exposed to elevated levels of circulating thyroid hormones, thyroxine (3,5,3',5'-tetraiodo- L-thyronine, or T_4) and triiodothyronine (3,5,3'-triiodo-L-thyronine, or T_3). Clinically, this state often manifests itself in weight loss, hypermetabolism, lowering of serum LDL levels, cardiac arrhythmias, heart failure, muscle weakness, bone loss in postmenopausal women, and anxiety. In most instances, thyrotoxicosis is due to hyperthyroidism, a term reserved for disorders characterized by overproduction of thyroid hormones by the thyroid gland. The ideal treatment of hyperthyroidism would be the elimination of its cause. This is however not possible in the more common diseases producing thyroid hypersecretion. At present, treatment of hyperthyroidism is directed to reduce overproduction of thyroid hormones by inhibiting their synthesis or release, or by ablating thyroid tissue with surgery or radioiodine.


Drugs inhibiting thyroid hormone synthesis, release or peripheral conversion of T_4 to T_3 include antithyroid drugs (thionamides), iodide, iodinated contrast agents, potassium perchlorate and glucocorticoids. The main action of antithyroid drugs such as methimazole (MMI), carbimazole, and propylthiouracil (PTU), is to inhibit the organification of iodide and coupling of iodotyrosines, thus blocking the synthesis of thyroid hormones. As they neither inhibit iodide transport or block the release of stored thyroid hormones, control of hyperthyroidism is not immediate and in most cases requires 2 to 6 weeks. Factors that determine the speed of restoration of euthyroidism include disease activity, initial levels of circulating thyroid hormones, and intrathyroidal hormone stores. Serious side effects are not common with antithyroid drugs. Agranulocytosis is the the most feared problem and has been

observed with both MMI or PTU treatment. The elderly may be more susceptible to this side effect, but agranulocytosis can occur in any age group, although less frequently. Inorganic iodide given in pharmacological doses (as Lugol's solution or as saturated solution of potassium iodide, SSKI) decreases its own transport into the thyroid, thus inhibiting iodide organification (the Wolff-Chaikoff effect), and rapidly blocks the release of T_4 and T_3 from the gland. However, after a few days or weeks, its antithyroid action is lost, and thyrotoxicosis recurs or may worsen. Short-term iodide therapy is used to prepare patients for surgery, usually in combination with a thionamide drug. Iodide is also used in the management of severe thyrotoxicosis (thyroid storm), because of its ability to inhibit thyroid hormone release acutely. Perchlorate interferes with accumulation of iodide by the thyroid. Gastric irritation and toxic reactions limit the long-term use of perchlorate in the management of hyperthyroidism. Glucocorticoids in high doses inhibit the peripheral conversion of T_4 to T_3 . In Graves' hyperthyroidism, glucocorticoids appear to decrease T_4 secretion by the thyroid, but the efficiency and duration of this effect is unknown. The aim of surgical treatment or radioiodine therapy of hyperthyroidism is to reduce the excessive secretion of thyroid hormones by removal or destruction of thyroid tissue. Subtotal or near-total thyroidectomy is performed in Graves' disease and toxic multinodular goiter. Restoration of euthyroidism before surgery is mandatory. The classical approach combines a course of thionamide treatment to restore and maintain euthyroidism, and the preoperative administration of iodide for approximately 10 days in order to induce involution of the gland. Propranolol and other beta-adrenergic antagonist drugs are useful in controlling tachycardia and other symptoms of sympathetic activation.

A high affinity ThR antagonist would in principle have the ability to restore euthyroidism quicker than any of the above agents, considered that its action is competitive for the ThR receptor. Such an agent could be used either alone or in combination with the above drugs, alternatively before an ablative treatment. It may also serve as a safer substitute for antithyroid drugs, especially in elderly patients at a high risk of agranulocytosis. Furthermore, hyperthyroidism can aggravate pre-existing heart disease and also lead to atrial fibrillation (AF), congestive heart failure, or worsening of angina pectoris. In an elderly patient, often with mild but prolonged elevation of plasma thyroid hormones, symptoms and signs of heart failure and complicating AF may dominate the clinical picture and mask the more classical endocrine manifestations of the disease.

DESCRIPTION OF THE INVENTION

In accordance with the present invention, compounds are provided which are thyroid receptor ligands, and have the general formula:



or a pharmaceutically acceptable salt thereof, wherein:

R₁ is selected from: C₃₋₁₀ aryl; C₃₋₁₀ heteroaryl; C₃₋₁₅ alkyl; C₄₋₁₅ alkenyl; C₃₋₁₅ alkynyl; and C₃₋₁₀ cycloalkyl, said aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl optionally being substituted with 1 to 3 groups of R^a;

R₂ is selected from: hydrogen; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; and C₁₋₄ alkoxy; or a bioisosteric equivalent;

R₁ may be linked through the available carbons to R₂, thus forming a C₄-C₈ cycloalkyl, saturated or partially unsaturated, and optionally substituted with 1 to 3 groups of R^a;

X is selected from: of -O-; -S-; -SO-; SO₂-; -Se-; -Te-; -N(R^c)-; and -S-S-;

R₃ and R₄ are the same or different and are selected from: halogen; C₁₋₄ alkyl; C₃₋₇ cycloalkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; -CF₃; -OCF₃; -OCF₂H; -SCH₃; -SCF₃; and -COOH, or a bioisosteric equivalent; with the proviso that when R₃ is C₁₋₄ alkoxy; -OCF₃; -OCF₂H; -SCH₃; -SCF₃; or -COOH, then R₄ must be selected from: halogen; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₃₋₇ cycloalkyl; and -CF₃;

n is an integer of 0, 1, 2 or 3;

Y is selected from: of -CO-; -O-; -S-; -CH(R^b)-; and -N(R^c)-;

R₃ is independently selected from:

carboxylic acid (-CO₂H);

phosphonic acid (-PO(OH)₂);

phosphamic acid (-PO(OH)NH₂);

sulphonic acid (-SO₂OH);

α-keto carboxylic acid (COCO₂H);

hydroxamic acid (-CONHOH);

sulphonamide (-SO₂NHR');

sulphonamide (-NH₂SO₂R'), with the proviso that then Y is CO and n is 0;

acylsulphonamide (-CONH₂SO₂R') or (-SO₂NHCOR');

or any other possible bioisosteric equivalent of all the groups above;

R^a represents a member selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; -SMe; -CN; -NO₂; and -N(C₀₋₄)₂, or a bioisosteric equivalent;

R^b represents a member selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; and -(CH₂)_mOH, where m is an integer between 0 and 4, or a bioisosteric equivalent;

R^c represents a member selected from: hydrogen; C₁₋₄ alkyl; C₃₋₄ alkenyl; and C₃₋₄ alkynyl or a bioisosteric equivalent;


included for the variables above are all the possible stereoisomers thereof;
prodrug ester forms thereof; and radioactive forms thereof.

DETAILED DESCRIPTION OF THE INVENTION

The following definitions apply to the terms as used throughout this specification, unless otherwise limited in specific instances.

The term "thyroid receptor ligand" as used herein is intended to cover any chemical substance which binds to a thyroid receptor. The ligand may act as an antagonist, an agonist, a partial antagonist or a partial agonist.

The term "alkyl" as employed herein alone or as part of another group refers to acyclic straight or branched chain radical, containing 1 to 15 carbons, preferably 1 to 10 carbons in the normal chain, i.e. methyl, ethyl, propyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl. Alkyl also refer to a radical where 1 to 6 hydrogens can be replaced by halogen through the available carbons. Alkyl also includes a straight or branched alkyl group which contains or is interrupted by a cycloalkane, preferable cyclopentane and cyclohexane, as exemplified below:



Wherein: when w and z is = 1 to 9, the sum of w and z is not more than 10; and when x and y is = 1 to 8, the sum of x and y is not more than 9.

The alkyl portions can be attached at any variable point of attachment to the cycloalkane. When a substituted alkyl is present, this refers to a straight or branched alkyl group as defined above, substituted with 1-3 groups of R^a as defined with respect to each variable.

The term "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 15 carbons and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to 5 carbon to carbon bonds may be present. Preferably 2 to 10 carbons are present in the normal chain radical, such as ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, decenyl, dodecenyl and the like. As described above with respect to the "alkyl", the straight or branched portion of the alkenyl group may be optionally substituted when a substituted alkenyl group is provided.

The term "alkynyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 15 carbons and at least one carbon to carbon triple bond. Preferably one carbon to carbon triple bond is present, and up to 5 carbon to carbon triple bonds may be present. Preferably 2 to 10 carbons are present in the normal chain, such as ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, decynyl, dodecynyl and the like. As described above with respect to the "alkyl", the straight or branched portion of the alkynyl group may be optionally substituted when a substituted alkynyl group is provided.

The term "cycloalkyl" as employed herein alone or as part of another group refers to saturated cyclic hydrocarbon groups or partially unsaturated cyclic hydrocarbon groups, independently containing 1 to 2 carbon to carbon double bonds or carbon to carbon triple bonds. The cyclic hydrocarbon contain 3 to 10 carbons, including rings that are fused. It should also be understood that the present invention involves cycloalkyl rings where 1 to 2 carbons in the ring are replaced by either -O-, -S- or -N-, thus forming a saturated or partially saturated heterocycle. Preferred cycloalkyl groups include 3 to 6 carbons, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, which may be optionally substituted through available carbons as in the case of "alkyl".


The term "aryl" as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups, consisting of 5 to 10 carbons in the ring portion, including partially saturated rings such as indanyl and tetrahydronaphthyl. The preferred aryl groups are phenyl and naphthyl, which may be substituted with 0-3 groups selected from R^a.

The term "halogen" or "halo" as used herein alone or as part of another group, exemplified by "haloalkoxy", refers to chlorine, bromine, fluorine and iodine.


The term "alkoxy" refers to those groups of the designated carbon length in either a straight or branched configuration attached through an oxygen linkage and if two or more carbons are in the length, they may include a double or a triple bond. Examples of such alkoxy groups are methoxy, ethoxy, propoxy, allyloxy, propargyloxy, butoxy, isobutoxy, tertiary butoxy, and the like. Alkoxy also refers to a radical where 1 to 3 hydrogens can be replaced by halogen through the available carbons, where the preferred group radical is -OCF₃.

The term "lower alkyl", "lower alkenyl" and "lower alkynyl" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons, containing 1 to 4 carbons, such as methyl, ethyl, vinyl, acetylene, n-propyl, i-propyl, butyl, t-butyl and the like. Lower alkyl also refers to a radical where 1 to 3 hydrogens can be replaced by halogen through the available carbons, where the preferred group radical is $-\text{CF}_3$.

The term "heteroaryl" or "heteroaromatic moiety" as used herein alone or as a part of another group refers to a group containing 5 to 10 atoms, where the aromatic ring includes 1 to 4 heteroatoms, such as nitrogen, oxygen or sulfur. Such rings may be fused to another aryl or heteroaryl ring, and includes possible N-oxides. The heteroaryl group may optionally be substituted by the available carbons with 0 to 3 substituents of R^a and also include substitution with carbonyl, hydroxy, and/or carboxyl. When R_5 is selected from heterocycles it refers to mainly to 5 membered rings, containing at least one nitrogen, as exemplified and depicted below:



The term "phosphonic acid" refers to a phosphorus containing group of the structure:



Wherein R and R' are independently selected from H or lower alkyl.

The term "bioisosteric equivalent" refers to compounds or groups that possess near equal molecular shapes and volumes, approximately the same distribution of electrons, and which

exhibit similar physical and biological properties. Examples of such equivalents are: (i) fluorine vs. hydrogen, (ii) oxo vs. thia, (iii) hydroxyl vs. amide, (iv) carbonyl vs. oxime, (v) carboxylate vs. tetrazole. Examples of such bioisosteric replacements can be found in the literature and examples of such are given below:

- 1) Burger, A. *Relation of chemical structure and biological activity*; in *Medicinal Chemistry* Third ed.; Burger, A., ed.; Wiley-Interscience: New York, 1970, pp. 64-80.
- 2) Burger, A.; "Isosterism and bioisosterism in drug design"; *Prog. Drug Res.* **1991**, *37*, 287-371.
- 3) Burger, A.; "Isosterism and bioanalogy in drug design"; *Med. Chem. Res.* **1994**, *4*, 89-92.
- 4) Clark, R. D.; Ferguson, A. M.; Cramer, R. D.; "Bioisosterism and molecular diversity"; *Perspect. Drug Discovery Des.* **1998**, *9/10/11*, 213-224.
- 5) Koyanagi, T.; Haga, T.; "Bioisosterism in agrochemicals."; *ACS Symp. Ser.* **1995**, *584*, 15-24.
- 6) Kubinyi, H.; "Molecular similarities. Part 1. Chemical structure and biological activity"; *Pharm. Unserer Zeit* **1998**, *27*, 92-106.
- 7) Lipinski, C. A.; "Bioisosterism in drug design"; *Annu. Rep. Med. Chem.* **1986**, *21*, 283-91.
- 8) Patani, G. A.; LaVoie, E. J.; "Bioisosterism: A rational approach in drug design"; *Chem. Rev. (Washington, D. C.)* **1996**, *96*, 3147-3176.
- 9) Soskic, V.; Joksimovic, J.; "Bioisosteric approach in the design of new dopaminergic/serotonergic ligands"; *Curr. Med. Chem.* **1998**, *5*, 493-512.
- 10) Thornber, C. W.; "Isosterism and molecular modification in drug design"; *Chem. Soc. Rev.* **1979**, *8*, 563-80.

The compounds of formula I can be present as salts, in particular "pharmaceutically acceptable salts". A compound having at least one acid group (for example -COOH) can form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono, di or trilower alkylamine, for example ethyl, tertbutyl, diethyl, diisopropyl, triethyl, tributyl or dimethyl-propylamine, or a mono, di or trihydroxy lower alkylamine, for example mono, di or triethanolamine. Corresponding internal salts may furthermore be formed. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds of formula I or their pharmaceutically

acceptable salts, are also included. Preferred salts of the compounds of formula I include sodium, potassium and magnesium salts and pharmaceutically acceptable organic amines.

An acid center (for example -COOH) part in formula I can form "prodrug ester forms" known in the art such as pivaloyloxymethyl or dioxolenylmethyl. Such prodrug esters are described in standard references such as Chapter 31, written by Camille G. Wermuth et al., in "The Practice of Medicinal Chemistry", ed. C. G. Wermuth, Academic Press, 1996 (and the references contained therein).

Certain compounds of the invention can be "stereoisomers", which have one or more asymmetric centers and can exist in the form of racemates, single enantiomers, as individual diastereomers, with all possible isomers, and mixtures thereof, all of which are within the scope of the invention.

Preferred salts of the compounds of formula I include sodium, potassium and magnesium salts and pharmaceutically acceptable organic amines.

A preferred embodiment of formula I is when X is -O-.

Another preferred embodiment of formula I is when R₂ is selected from hydrogen and methyl; X is -O-; R₃ and R₄ are the same or different and are halogen, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, or -CF₃; n is an integer of 0 or 1; Y is -CH(R^b)-; R₅ is carboxylic acid (-CO₂H); R^b is hydrogen.

Yet another preferred embodiment of formula I is when X is -O-; Y is -N(R^c)-; n is an integer of 0, 1 and 2; R₅ is a carboxylic acid (-CO₂H) or an α-keto carboxylic acid (COCO₂H); R^c is hydrogen.

Yet another preferred embodiment of formula I is when X is -O-; Y is -O-; n is an integer of 1 and 2; R₅ is a carboxylic acid (-CO₂H).


Yet another preferred embodiment of formula I is when X is -O-; Y is -CO-; n is 0; R₅ is a sulphonamide (-NHSO₂R').














Yet another preferred embodiment of formula I is when R_2 is selected from hydrogen and methyl; X is -O-; R_3 is -COOH and R_4 is selected from: halogen; C_{1-4} alkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; and -CF₃; R^b is hydrogen; R^c is hydrogen.

The compounds of formula I may be prepared by the exemplary processes described in the following reaction schemes. Exemplary reagents and procedures for these reactions appear hereinafter and in the working Examples.












Compounds of formula I of the invention can be prepared using either method A, B or C as indicated in Scheme 1 below. In Scheme 1 (Method A, B or C) the phenol **1** is dissolved in a suitable solvent, treated with 1-5 molar equivalents of an appropriate base, such as potassium carbonate, cesium carbonate, potassium hydroxide, sodium hydride or triethylamine. The resulting anion is then O-alkylated with the appropriate halides R-Br, R-Cl or R-OMs. Other combinations of alkylating agents or bases may be employed and are known to those skilled in the art. The reaction mixture is stirred at room temperature or heated until the starting materials are consumed. After standard work-up, the ester function is removed by treatment with 3-6 molar equivalents of a base such as sodium hydroxide at room temperature, dissolved in a solvent such as methanol. The reaction mixture gives after acidification with hydrochloric acid a precipitate which is the end product **2** (Examples 1-20). In Examples 21-30 the end products **2** are further purified by column chromatography.







Scheme 1



Example	R	n
1		1
2		1
3		1
4		1
5		1
6		1
7		1
8		2
9		2
10		2
11		2
12		2
13		2

14

- | | | |
|----|---|---|
| 14 |  | 2 |
| 15 |  | 2 |
| 16 |  | 2 |
| 17 |  | 2 |
| 18 |  | 2 |
| 19 |  | 2 |
| 20 |  | 1 |
| 21 |  | 1 |
| 22 |  | 2 |
| 23 |  | 2 |
| 24 |  | 2 |

25		2
26		2
27		2
28		2
29		2
30		2


Examples of compounds of formula I in which further variation at the R₁, R₃, R₄ and Y positions is introduced are shown in Scheme 2. A method similar as above is used for the O-alkylation step (Method D), but the preparations may be done in an automated fashion (Examples 36-100). After standard work-up, the ester function can be hydrolysed by 4 different methods (D1-D4) to give the end-products 4.




In the first method (D1), the ester is dissolved in methanol, sodium hydroxide is added and the mixture stirred at room temperature. After acidification with hydrochloric acid the product is extracted into ethyl acetate. In the second method (D2), the ester is dissolved in dry dichloromethane, potassium trimethylsilanolate is added and the reaction mixture is stirred at room temperature. After concentration, the residue is re-dissolved in methanol and











neutralised on an SCX SPE column. In the third method (D3), the ester is dissolved in dry tetrahydrofuran, potassium trimethylsilanolate is added and the reaction mixture stirred at room temperature. The mixture is neutralised on an SCX SPE column. In the last method (D4), the ester is dissolved in tetrahydrofuran, lithium hydroxide is added and the reaction mixture stirred at room temperature. The mixture is neutralised on an SCX SPE column. In all methods, the final products are further purified on a silica SPE column.












Several other related methodologies exist for the hydrolysis of alkyl esters and are known to those skilled in the art. Furthermore, other protecting groups for the carboxylic acid can be employed, and their usage is known to those skilled in the art (references describing protecting group strategy include, for example, "Protecting Groups in Organic Chemistry", J. F. W. McOmie, Plenum Press, London, New York, 1973, and "Protective Groups in Organic Synthesis", T. W. Greene, Wiley, New York, 1984).












Scheme 2






















Example	R	R ₃ ,R ₄	Y	n
36		Br	CH ₂	0
37		Br	CH ₂	0
38		Br	CH ₂	1










39		Br	CH ₂	1
40		Br	CH ₂	1
41		Br	CH ₂	1
42		Cl	CHOH	0
43		Cl	CHOH	0
44		Cl	CHOH	0
45		Cl	C=O	0
46		Cl	C=O	0
47		Cl	C=O	0
48		Cl	C=O	0


49		Cl	C=O	0
50		Me	CH ₂	0
51		Me	CH ₂	0
52		Me	CH ₂	0
53		Me	CH ₂	0
54		Me	CH ₂	0
55		Me	CH ₂	0
56		Me	C=O	0
57		Me	C=O	0
58		Me	C=O	0
59		Me	C=O	0

60		Me	C=O	0
61		<i>i</i> -Pr	C=O	0
62		<i>i</i> -Pr	C=O	0
63		<i>i</i> -Pr	C=O	0
64		<i>i</i> -Pr	C=O	0
65		<i>i</i> -Pr	C=O	0
66		<i>i</i> -Pr	C=O	0
67		<i>i</i> -Pr	C=O	0
68		<i>i</i> -Pr	CH ₂	0
69		<i>i</i> -Pr	CH ₂	0
70		<i>i</i> -Pr	CH ₂	0

71		<i>i</i> -Pr	CH ₂	0
72		Br	CH ₂	1
73		Br	CH ₂	1
74		Br	CH ₂	1
75		Br	CH ₂	1
76		Br	CH ₂	1
77		Br	CH ₂	1
78		Br	CH ₂	1
79		Br	CH ₂	1
80		Br	CH ₂	1


81		Br	CH ₂	1
82		Br	CH ₂	1
83		Br	CH ₂	1
84		Br	CH ₂	1
85		Br	CH ₂	1
86		Br	CH ₂	1
87		Br	CH ₂	1
88		Br	CH ₂	1
89		Br	CH ₂	1





Further variations of the R₃ and R₄ groups are shown in Scheme 3, where R₄ is equal to a methoxy group and R₃ is equal to a bromine. In the referred Examples 109 and 110 the same coupling procedures as above were employed.

Scheme 3




Scheme 4 depicts a synthesis of compounds of formula I in which -Y-(CH₂)_n-R₅ is an acylsulphonamide (when Y = CO, n = 0 and R₅ = -NHSO₂R).

In one procedure, 8 is mixed with a base such as N,N-diisopropylethylamine (DIEA) and the appropriate sulphonamide in a mixture of dichloromethane and dimethylformamide.

Treatment of the mixture with a base and coupling reagent combinations such as benzotriazole-1-yl-oxy-tris-pyrrolidino phosphonium hexafluorophosphate (PyBOP) and 1-hydroxybenzotriazole hydrate (HOBt), gives after stirring at room temperature and subsequent mild acid treatment during work-up and purification by HPLC, the desired acylsulphonamides 9 (Example 31-33, 35).

Several other procedures for the preparation of acylsulphonamides can be employed. For example, a carboxylic acid, a coupling reagent such as 3-ethyl-1-[3-(dimethylamino)propyl]carbodiimide hydrochloride (EDCI), and a base such as dimethylaminopyridine (DMAP) and a sulphonamide in dichloromethane can be stirred at room temperature, to give an acylsulphonamide.

Scheme 4




Example	R	R'
31		
32		
33		
35		Me

Additional examples of transformations of the Examples is shown in Scheme 5, where the nitro group of Example 33 is reduced to an amino group to give 10 (Example 34) by


employing sodium dithionite. Several other reducing agents are known to those skilled in the art and the referred example should not be considered as limiting.

Scheme 5




Yet further examples of variations of the $-Y-$, $-(CH_2)_n-$ and $-R_5$ groups is shown in Scheme 6, where $-Y-(CH_2)_n-R_5$ is an oxamic acid (when $Y=NH$, $n=0$ and $R_5=COCO_2H$). A nitro compound 11 group is reduced to an amino group, which subsequently is coupled with diethyl oxalate at elevated temperatures to give after standard ester saponification, the desired oxamic acid derivative 12 (Examples 107 and 108).

Scheme 6




Examples where two O-alkylations are done in a consecutive order is depicted in Scheme 7. Following similar procedures as described above (Method D), methyl 3-(3,5-dibromo-4-hydroxyphenyl)propionate is reacted with 3-bromomethyl- phenyl acetate, hydrolyzed to give the intermediate 13 and then further reacted with the exemplified lower haloalkyls, to give 14 (Examples 101, 102 and 103). Other alternative combinations of starting phenols, halides and procedures can be employed and are obvious for those skilled in the art.

Scheme 7



A similar strategy as above is exemplified and depicted in Scheme 8. Amino compound 15 (Example 104) can be further reacted with acetic acid under reductive conditions at different temperatures. Depending on temperature and reaction time, the mono or di-N-alkylated product is obtained. Many alternative procedures for the N-alkylation of aromatic amines can be employed and are obvious for those well skilled in the art.


Scheme 8



With respect to the above reaction scheme, although the various R_1 , R_2 , R_3 , R_4 , R_5 , X, Y and n moieties are specifically defined, unless otherwise indicated, it is to be understood that R_1 ,

R_2 , R_3 , R_4 , R_5 , X and Y may be any of the groups encompassed thereby and n may be 0, 1, 2, or 3.

Another aspect of the present invention refers to the use of a compound in medical therapy according to the formula below:



wherein:

R_1 is selected from: of C_{3-10} aryl; C_{5-10} heteroaryl; C_{3-15} alkyl; C_{4-15} alkenyl; C_{3-15} alkynyl; and C_{3-10} cycloalkyl, said aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl optionally being substituted with 1 to 3 groups of R^a ;

R_2 is selected from: hydrogen; C_{1-4} alkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; and C_{1-4} alkoxy, or a bioisosteric equivalent;

R_1 may be linked through the available carbons to R_2 , thus forming a C_4 - C_8 cycloalkyl, saturated or partially unsaturated, and optionally substituted with 1 to 3 groups of R^a ;

X is selected from: -O-; -S-; -SO-; SO_2 -; -Se-; -Te-; -N(R^c)-; and -S-S-;

R_3 and R_4 are the same or different and are selected from: halogen; C_{1-4} alkyl; C_{3-7} cycloalkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; C_{1-4} alkoxy; -CF₃; -OCF₃; -OCF₂H; -SCH₃; -SCF₃; -COOH or a bioisosteric equivalent;

Y is selected from: -CO-; -O-; -S-; -CH(R^b)-; and -N(R^c)-;

R_5 is independently selected from carboxylic acid (-CO₂H); phosphonic acid (-PO(OH)₂); phosphamic acid (-PO(OH)NH₂); sulphonic acid (-SO₂OH); α -keto carboxylic acid (COCO₂H); hydroxamic acid (-CONHOH); sulphonamide (-SO₂NHR'); sulphonamide

(-NHSO₂R'); acylsulphonamide (-CONHSO₂R') or (-SO₂NHCOR') or any other possible bioisosteric equivalent of all the groups above;

R₆ and R₇ are the same or different and are selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; and -(CH₂)_mOH; where m is an integer between 1 and 4, or a bioisosteric equivalent;

R^a is selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; -SMe; -CN; -NO₂; and -N(C₀₋₄)₂, or a bioisosteric equivalent;

R^b represents a member selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; and -(CH₂)_mOH, where m is an integer between 0 and 4, or a bioisosteric equivalent;

R^c is selected from: hydrogen; C₁₋₄ alkyl; C₃₋₄ alkenyl; and C₃₋₄ alkynyl, or a bioisosteric equivalent;

or pharmaceutically acceptable salts thereof as well as included for the variables above are all the possible stereoisomers thereof; prodrug ester forms thereof; and radioactive forms thereof.

A preferred embodiment of this aspect of the present invention are the use of the compounds referred to above wherein:

R₂ is selected from hydrogen and methyl;


X is -O-;








R₃ and R₄ are the same or different and are selected from: halogen; C₁₋₄ alkoxy; C₁₋₄ alkyl; and -CF₃;


R₅ is a carboxylic acid (-CO₂H);

R₆ and R₇ is hydrogen.

A particularly preferred embodiment of this aspect of the present invention relates to, but not limited to the use of the following compounds :



R	R ₄
	Br
	Br
	Br
	Br
	Br
	Br
	Br



or pharmaceutically acceptable salts thereof and stereoisomers thereof.

The compounds of the invention are antagonists or partial antagonists, preferably α -selective. As such they are useful in medical therapy. Furthermore, they are useful in the prevention, inhibition or treatment of a disease which is dependent on the expression of a T_3 regulated gene or associated with metabolic dysfunction. Examples of such diseases are heart related disorders, such as cardiac arrhythmias (atrial and ventricular arrhythmias), especially atrial fibrillation and ventricular tachycardia and fibrillation. The compounds of the invention may also be useful for the treatment of thyrotoxicosis, especially in the therapy of elderly patients, subclinical hyperthyroidism, and other related endocrine disorders, related to thyroid hormone.

The compounds of the invention may also be used to treat certain skin disorders or diseases such as keloids, roughened skin, lichen planus, ichthyosis, acne, psoriasis, Dernier's disease, eczema, chloracne, atopic dermatitis, pityriasis, hirsutism and skin scarring. In treating skin disorders or diseases as described above, the compounds of the invention may be used in combination with a retinoid or a vitamin D analog.

Exemplifying the invention is a pharmaceutical composition comprising any of the compounds described above and a pharmaceutically acceptable carrier. Also exemplifying the invention is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

Another embodiment of the invention is a method of treating, inhibiting or preventing a disease which is dependent on the expression of a T_3 regulated gene or associated with metabolic dysfunction in a mammal in need thereof by administering to the mammal a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above. The said diseases may be heart related disorders, such as cardiac arrhythmias (atrial and ventricular arrhythmias), especially atrial fibrillation and ventricular tachycardia and fibrillation, especially in the therapy of elderly patients, subclinical hyperthyroidism, and other related endocrine disorders, related to thyroid hormone.

Yet another embodiment of the invention is a method of treating, inhibiting or preventing certain skin disorders or diseases such as keloids, roughened skin, lichen planus, ichthyosis, acne, psoriasis, Dernier's disease, eczema, chloracne, atopic dermatitis, pityriasis, hirsutism and skin scarring. In treating skin disorders or diseases as described above, the compounds of the invention may be used in combination with a retinoid or a vitamin D analog.

Further exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment, inhibition or prevention of a disease which is dependent on the expression of a T_3 regulated gene or associated with metabolic dysfunction. Still further exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment and/or prevention of heart related disorders, such as cardiac arrhythmias (atrial and ventricular arrhythmias), especially atrial fibrillation and ventricular tachycardia and fibrillation, especially in the therapy of elderly patients, subclinical hyperthyroidism, and other related endocrine disorders, related to thyroid hormone.

Further exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment, inhibition or prevention of certain skin disorders or diseases such as keloids, roughened skin, lichen planus, ichthyosis, acne, psoriasis, Dernier's disease, eczema, chloracne, atopic dermatitis, pityriasis, hirsutism and skin scarring. In treating skin disorders or diseases as described above, the compounds of the invention may be used in combination with a retinoid or a vitamin D analog.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powder, granules, elixirs, tinctures, suspensions, syrups and emulsions. Likewise, they may also be administered in intravenous (bolus or infusion), intraperitoneal, topical (*e.g.*, ocular eyedrop), subcutaneous, intramuscular, or transdermal (*e.g.*, patch) form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex, and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician, veterinarian or clinician can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

Oral dosages of the present invention, when used for the indicated effects, will range between about 0.01 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 mg per kg of body weight per day (mg/kg/day) to 10 mg/kg/day, and most preferably 0.1 to 5.0 mg/kg/day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A medicament typically contains from about 0.01 mg to about 500 mg of the active ingredient, preferably from about 1 mg to about 100 mg of active ingredient. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg/minute

during a constant rate infusion. Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include without limitation starch, methylcellulose, agar, bentonite, xanthan gum and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example in "Design of Prodrugs" ed. H. Bundgaard, Elsevier, 1985, which is incorporated by reference herein in its entirety. Metabolites of the compounds includes active species produced upon introduction of compounds of this invention into the biological milieu.

The following Examples represent preferred embodiments of the present invention. However, they should not be construed as limiting the invention in any way. The ¹H NMR spectra was all consistent with the assigned structures in the Examples.

GENERAL PROCEDURE FOR THE PREPARATION OF EXAMPLES 1-13

(METHOD A).

A mixture of the appropriate phenol (1.5 mmol) and potassium carbonate (0.85 g, 6.2 mmol) in acetone (3 mL) was stirred at room temperature. After 10 minutes the bromo compound (6.2 mmol) was added. The reaction mixture was heated at reflux over night, cooled to room temperature and concentrated *in vacuo*. Diethyl ether was added to the residue, the solution filtered on silica gel and the resulting filtrate concentrated. The residue was dissolved in 12 mL of methanol. An aqueous solution of sodium hydroxide (8 mL, 1 N) was added dropwise and the reaction mixture was stirred at room temperature for 16 hours, and acidified with hydrochloric acid (1N). The precipitate formed was collected and dried.


GENERAL PROCEDURE FOR THE PREPARATION OF EXAMPLES 14-20(METHOD B).

A solution of the phenol (0.1 mmol), triethylamine (0.4 mmol) and acetone (0.5 mL) was stirred under nitrogen for 30 minutes. This solution was added to the appropriate bromide (0.4 mmol) in acetone (0.3 mL), the vessel sealed under nitrogen, and heated at 60 °C over night. After cooling to room temperature, the reaction mixture was concentrated *in vacuo*. The residue was suspended in sodium hydroxide (1N, 1 mL), the vessel sealed and stirred at room temperature. After 24 hours, hydrochloric acid (6 N, 0.5 mL) was added followed by ethyl acetate (2 mL). The mixture was shaken and placed on a separate ChemElute cartridge. After 5 minutes the cartridge was washed successively with ethyl acetate (3x1 mL), the precipitate collected and dried.












GENERAL PROCEDURE FOR THE PREPARATION OF EXAMPLES 21-30(METHOD C).








A mixture of the appropriate phenol (0.36 mmol) and potassium carbonate (0.2 g, 1.45 mmol) in 6 mL of acetone was stirred at room temperature. After 10 minutes the bromo compound (1.4 mmol) was added. The reaction mixture was heated under reflux over night, cooled to room temperature and concentrated *in vacuo*. Diethyl ether was added to the residue and filtered through a celite pad. The resulting filtrate was concentrated and the residue was dissolved in methanol (10 mL). An aqueous solution of sodium hydroxide (2 mL, 1N) was added dropwise and the reaction mixture was stirred over night at room temperature. The solution was acidified with hydrochloric acid (2 N) and the volatiles were removed *in vacuo*. The precipitate was collected, washed with water and purified by column chromatography (silica gel, gradient of chloroform and chloroform/methanol (from 90-10%).

For a table of Examples 1-30 comprising the Examples, methods, yields and mass spectras, see Scheme below.



Example	R	n	Method	Yield ¹	MS ²
1		1	A	89	365.2
2		1	A	32	365.2
3		1	A	81	379
4		1	A	90	379
5		1	A	71	393.1
6		1	A	79	393.1
7		1	A	74	405.1
8		2	A	68	379
9		2	A	88	293.1
10		2	A	92	407.2
11		2	A	90	393.1
12		2	A	70 ³	407.2

13		2	A	63	419.2
14		2	B	30	-
15	 Me	2	B	66	428.1
16		2	B	64	391
17		2	B	82	412.9
18		2	B	77	390.4
19		2	B	50	-
20		1	A	96	365.2
21		1	C	37	400.1
22	 CF ₃	2	C	84	482.1
23	 Br	2	C	67	493

24		2	C	97	428.1
25		2	C	95	428.1
26		2	C	75	442.1
27		2	C ⁴	5	432.1
28		2	C ⁴	42	482.1
29		2	C	65	459.1
30		2	C ⁴	4	470.2

¹⁾ Yields in %.

²⁾ MS result obtained on a Perkin-Elmer API 150Ex spectrometer, using electrospray negative ion mode.

³⁾ This substance fell as an oil during final work-up. In order to get crystals the methanol was removed in vacuo and the remaining water phase, extracted with ethyl acetate. After drying over magnesium sulphate and removal of the organic phase, a crystal mass was obtained.

⁴⁾ In Examples 27, 28 and 30, the final compound was first separated in 2 g silica pre-packed in 3 mL SPE cartridges employing the same gradient of solvents described in Procedure C and then further purified by semi-preparative-HPLC (Zorbax CombiHT (SB-C8 50x21.2 mm, 5μ) Mobil Phase: Solvent A. Water with 0.5% formic acid; Solvent B: acetonitrile. Gradient: 2 min 80% of A then over 8 min to 5% of A).

Example 31***N*-[3,5-Dibromo-4-(3-bromobenzyloxy)benzoyl]benzenesulphonamide**

To a stirred mixture of benzenesulphonamide (0.89 mmol), dichloromethane (1 mL) and dimethylformamide (one drop), a solution of 3,5-dibromo-4-(3-bromobenzyloxy)benzoic acid (0.17 mmol) and *N,N*-diisopropylethylamine (0.86 mmol) in dichloromethane (1 mL) was added. After 15 minutes bromotripyrrolidino phosphoniumhexafluoro phosphate (0.21 mmol) and 1-hydroxybenzotriazole hydrate (0.045 mmol) were added. The reaction mixture was stirred at room temperature over night. Dichloromethane was added (3 mL) and the mixture was washed with an aqueous solution of citric acid (10 %, 3x3 mL). The organic layer was washed with brine, dried over magnesium sulphate and concentrated *in vacuo*. The residue was purified on column (silica gel, chloroform/methanol 19:1) and further purified on semi-preparative HPLC (column and eluents as described for Examples 27, 28 and 30) to give 16 mg (16 %) of *N*-[3,5-dibromo-4-(3-bromobenzyloxy)benzoyl]benzenesulphonamide.

Example 32***N*-[3,5-Dibromo-4-(3-bromobenzyloxy)benzoyl]-3-nitrobenzenesulphonamide**

3,5-dibromo-4-(3-bromobenzyloxy)benzoic acid (0.17 mmol) was coupled with 3-nitrobenzenesulphonamide (0.89 mmol) using the procedure described in Example 31.

Purification on column (silica gel, chloroform/methanol 9:1) gave 60 mg (54%) of *N*-[3,5-dibromo-4-(3-bromobenzyloxy)benzoyl]-3-nitrobenzenesulphonamide.

Example 33***N*-[3,5-Dibromo-4-(3-bromobenzyloxy)benzoyl]-4-nitrobenzenesulphonamide**

3,5-Dibromo-4-(3-bromobenzyloxy)benzoic acid (0.13 mmol) was coupled with 4-nitrobenzenesulphonamide (0.64 mmol) using the procedure described in Example 31.

Purification on column (silica gel, chloroform/methanol 9:1) gave 58 mg (69%) of *N*-[3,5-dibromo-4-(3-bromobenzyloxy)benzoyl]-4-nitrobenzenesulphonamide.

Example 34**4-Amino-*N*-[3,5-dibromo-4-(3-bromobenzyloxy)benzoyl]benzenesulphonamide**

To a solution of *N*-[3,5-dibromo-4-(3-bromobenzyloxy)benzoyl]-4-nitro-benzenesulphonamide (0.25 mmol), prepared as described in Example 33, in ethanol (95 %, 25 mL), sodium dithionite (1.26 mmol) was added and the reaction mixture stirred over night at 70°C. The reaction mixture was allowed to reach room temperature and then evaporated *in vacuo*. The residue was diluted with ethyl acetate and washed with an aqueous solution of sodium hydrogencarbonate (saturated). The organic phase was dried over magnesium sulphate and concentrated *in vacuo*. The residue was purified on column (silica gel, chloroform/methanol 17:3) and further purified by semi-preparative HPLC (column and eluents as described for Examples 27, 28 and 30) to give 30 mg (20 %) of 4-amino-*N*-[3,5-dibromo-4-(3-bromobenzyloxy)-benzoyl]benzenesulphonamide.

Example 35

***N*-[3,5-Dibromo-4-(3-bromobenzyloxy)benzoyl]methanesulphonamide**

3,5-Dibromo-4-(3-bromobenzyloxy)benzoic acid (0.15 mmol) was coupled with methanesulphonamide (0.77 mmol) using the procedure described in Example 31. Purification on column (silica gel, chloroform/methanol 9:1) gave 55 mg (67%) of *N*-[3,5-dibromo-4-(3-bromobenzyloxy)benzoyl]methanesulphonamide.

GENERAL PROCEDURE FOR THE PREPARATION OF EXAMPLES 36-100

(METHOD D1, D2, D3 AND D4)

To a solution of the appropriate phenol (0.035 mmol) in acetone or dry acetonitrile (0.25 mL) was added potassium carbonate (0.070 or 0.14 mmol) and the resulting mixture was stirred at room temperature. After 10 minutes a solution of the halide (0.070 or 0.14 mmol) in acetone or acetonitrile (0.25 mL) was added. When the halide was a chloride, a catalytic amount of sodium iodide was added. The mixture was heated at 60 °C (acetone) or 80°C (acetonitrile) over night. After cooling down to room temperature, the reaction mixture was filtered through a silica SPE column (500 mg/3 mL), eluting with *n*-heptane/ethyl acetate 3:1 (3 mL). After concentration *in vacuo* the residue was hydrolysed according to method D1, D2, D3 or D4.

Method D1: The residue was dissolved in methanol (0.25 mL), sodium hydroxide (1 N in water, 0.25 mL) was added and the mixture was stirred at room temperature over night. After acidification with hydrochloric acid (1 N) the product was extracted into ethyl acetate.


Method D2: The residue was dissolved in dry dichloromethane (0.5 mL) and potassium trimethylsilanolate (9 mg, 0.070 mmol, 2 equiv) was added. The reaction mixture was stirred at room temperature over night. After concentration, the residue was dissolved in methanol and neutralised on an SCX SPE column, using methanol as eluent.












Method D3: The residue was dissolved in dry tetrahydrofuran (0.5 mL) and potassium trimethylsilanolate (9 mg, 0.070 mmol, 2 equiv) was added. The reaction mixture was stirred at room temperature until TLC showed complete consumption of the starting material (0.5 – 2 h). The mixture was neutralised on an SCX SPE column, using methanol as eluent.












Method D4: The residue was dissolved in tetrahydrofuran (0.25 mL) and lithium hydroxide (1 N in water, 0.25 mL) was added. The reaction mixture was stirred at room temperature. After 4 hours the mixture was neutralised on an SCX SPE column (500 mg/3 mL), using methanol or triethylamine (10% in methanol) as eluent.












Methods D1-D4: The solvent was concentrated and the residue purified on a silica SPE column (500 mg/3 mL, heptane/ethyl acetate 9:1 (3 mL) followed by dichloromethane/methanol 9:1 (3 mL)). The product containing fractions were collected and concentrated *in vacuo* to give the final product.










For a table of Examples 36-100 comprising the Examples, used solvents, hydrolysis method, yields and mass spectras, see Scheme below.






















Example	R	R ₃ ,R ₄	Y	n	Solvent	Equiv. halide	Hydrol. Method	Yield (%)	MS ¹
36		Br	CH ₂	0	acetone	4	D1	27	421.0
37		Br	CH ₂	0	acetone	4	D4	75	363.1
38		Br	CH ₂	1	acetone	4	D2	34	463.0
39		Br	CH ₂	1	CH ₃ CN	2	D3	38	570.7
40		Br	CH ₂	1	CH ₃ CN	2	D3	51	437.8
41		Br	CH ₂	1	CH ₃ CN	2	D3	91	443.2
42		Cl	CHOH	0	CH ₃ CN	2	D4	58	290.8
43		Cl	CHOH	0	CH ₃ CN	2	D4	52	333.1
44		Cl	CHOH	0	CH ₃ CN	2	D4	64	404.8
45		Cl	C=O	0	CH ₃ CN	2	D4	48	289.0
46		Cl	C=O	0	CH ₃ CN	2	D4	12	330.7





47		Cl	C=O	0	CH ₃ CN	2	D4	64	402.7
48		Cl	C=O	0	CH ₃ CN	2	D4	7	351.1
49		Cl	C=O	0	CH ₃ CN	2	D4	63	352.9
50		Me	CH ₂	0	CH ₃ CN	2	D4	61	235.0
51		Me	CH ₂	0	CH ₃ CN	2	D4	46	275.2
52		Me	CH ₂	0	CH ₃ CN	2	D4	61	277.0
53		Me	CH ₂	0	CH ₃ CN	2	D4	78	349.0
54		Me	CH ₂	0	CH ₃ CN	2	D4	76	297.1
55		Me	CH ₂	0	CH ₃ CN	2	D4	85	299.2
56		Me	C=O	0	CH ₃ CN	2	D4	43	248.8
57		Me	C=O	0	CH ₃ CN	2	D4	82	290.8

58		Me	C=O	0	CH ₃ CN	2	D4	93	361.0
59		Me	C=O	0	CH ₃ CN	2	D4	85	310.9
60		Me	C=O	0	CH ₃ CN	2	D4	86	313.0
61		<i>i</i> -Pr	C=O	0	CH ₃ CN	2	D4	55	305.2
62		<i>i</i> -Pr	C=O	0	acetone	4	D1	26	361.3
63		<i>i</i> -Pr	C=O	0	CH ₃ CN	2	D4	53	344.8
64		<i>i</i> -Pr	C=O	0	CH ₃ CN	2	D4	51	346.9
65		<i>i</i> -Pr	C=O	0	CH ₃ CN	2	D4	61	417.1
66		<i>i</i> -Pr	C=O	0	CH ₃ CN	2	D4	72	367.0
67		<i>i</i> -Pr	C=O	0	CH ₃ CN	2	D4	72	368.8
68		<i>i</i> -Pr	CH ₂	0	CH ₃ CN	2	D4	41	291.1

69		<i>i</i> -Pr	CH ₂	0	CH ₃ CN	2	D4	25	405.1
70		<i>i</i> -Pr	CH ₂	0	CH ₃ CN	2	D4	23	353.2
71		<i>i</i> -Pr	CH ₂	0	CH ₃ CN	2	D4	25	355.0
72		Br	CH ₂	1	CH ₃ CN	2	D4	100	497.2
73		Br	CH ₂	1	CH ₃ CN	2	D4	86	499
74		Br	CH ₂	1	CH ₃ CN	2	D4	100	430.9
75		Br	CH ₂	1	CH ₃ CN	2	D4	100	448.9
76		Br	CH ₂	1	CH ₃ CN	2	D4	61	549.1
77		Br	CH ₂	1	CH ₃ CN	2	D4	91	478.9

78		Br	CH ₂	1	CH ₃ CN	2	D4	100	463.0
79		Br	CH ₂	1	CH ₃ CN	2	D4	49	413.8
80		Br	CH ₂	1	CH ₃ CN	2	D4	32	413.8
81		Br	CH ₂	1	CH ₃ CN	2	D4	58	413.8
82		Br	CH ₂	1	CH ₃ CN	2	D4	93	463.9
83		Br	CH ₂	1	CH ₃ CN	2	D4	9	502.9
84		Br	CH ₂	1	CH ₃ CN	2	D4	63	565.9
85		Br	CH ₂	1	CH ₃ CN	2	D4	88	418.0
86		Br	CH ₂	1	CH ₃ CN	2	D4	46	480.7
87		Br	CH ₂	1	CH ₃ CN	2	D4	38	430.9

88		Br	CH ₂	1	CH ₃ CN	2	D4	76	454.9
89		Br	CH ₂	1	CH ₃ CN	2	D4	46	433.9
90		Br	CH ₂	1	CH ₃ CN	2	D4	94	446.8
91		Br	CH ₂	1	CH ₃ CN	2	D4	97	446.8
92		Br	CH ₂	1	CH ₃ CN	2	D4	30	538.9
93		Br	CH ₂	1	CH ₃ CN	2	D4	76	421.0
94		Br	CH ₂	1	CH ₃ CN	2	D4	36	457.9
95		Br	CH ₂	1	CH ₃ CN	2	D4	17	478.9
96		Br	CH ₂	1	CH ₃ CN	2	D4	28	497.2

97		Br	CH ₂	1	CH ₃ CN	2	D4	29	559.0
98		Br	CH ₂	1	CH ₃ CN	2	D4	20	481.0
99		Br	CH ₂	1	CH ₃ CN	2	D4	25	541.0
100		Br	CH ₂	1	CH ₃ CN	2	D4	41	493.0

¹⁾ MS result obtained on a Perkin-Elmer API 150Ex spectrometer, using electrospray negative ion mode.

Example 101

3-[3,5-Dibromo-4-(3-ethoxybenzyloxy)phenyl]propionic acid

(a) Methyl 3-(3,5-dibromo-4-hydroxyphenyl)propionate (0.15 g, 0.44 mmol) was coupled with 3-bromomethylphenyl acetate (0.20 g, 0.88 mmol) using the procedure described in Method D4. Purification on column (silica gel, dichloromethane/methanol, gradient elution from 95:5 to 90:10) gave 123 mg (78%) of

3-[3,5-dibromo-4-(3-hydroxybenzyloxy)phenyl]propionic acid. MS: m/z 430.1 (M⁺-1).

(b) 3,5-dibromo-4-(3-hydroxybenzyloxy)phenyl]propionic acid was coupled with ethylbromide (4 equivalents) in acetonitrile, using the procedure described in Method D4, to give 9 mg (56%) of 3,5-dibromo-4-(3-ethoxybenzyloxy)phenyl]- propionic acid. MS: m/z 457.0 (M⁺-1).

Example 102**3-[3,5-Dibromo-4-(3-propyloxybenzyloxy)phenyl]propionic acid**

3-[3,5-Dibromo-4-(3-hydroxybenzyloxy)phenyl]propionic acid was coupled with propylbromide using the procedure described in Example 101(b), to give 8 mg (50%) of 3,5-dibromo-4-(3-propyloxybenzyloxy)phenyl]propionic acid. MS: m/z 471.1 (M^+-1).

Example 103**3-[3,5-Dibromo-4-(3-butyloxybenzyloxy)phenyl]propionic acid**

3,5-dibromo-4-(3-hydroxybenzyloxy)phenyl]propionic acid was coupled with butylbromide using the procedure described in Example 101(b), to give 8 mg (50%) of 3,5-dibromo-4-(3-butyloxybenzyloxy)phenyl]propionic acid. MS: m/z 484.9 (M^+-1)

Example 104**3-[3,5-Dibromo-4-(3-aminobenzyloxy)phenyl]propionic acid**

A mixture of 3-[3,5-dibromo-4-(3-nitrobenzyloxy)phenyl]propionic acid (0.66 g, 1.44 mmol) and $Na_2S_2O_4$ (2.0 g, 11.5 mmol) in ethanol (50 mL, 90%) was stirred at 70°C over night. The reaction mixture was concentrated and purified on column (silica gel, chloroform/methanol, 9:1), to give 0.18 g (29%) of 3-[3,5-dibromo-4-(3-amino- benzyloxy)phenyl]propionic acid. MS: m/z 429.1 (M^+-1).

Example 105**3-[3,5-Dibromo-4-(3-ethylaminebenzyloxy)phenyl]propionic acid**

Sodium boron hydride (6.6 mg, 0.17 mmol) was added in portions to a stirred solution of 3-[3,5-dibromo-4-(3-aminobenzyloxy)phenyl]propionic acid (15 mg, 0.035 mmol) in glacial acetic acid (2 mL) at room temperature. After 5.5 hours the reaction was quenched with water, extracted with diethyl ether and purified on semi-preperative HPLC using the same conditions as described for Examples 27, 28 and 30. This gave 4.0 mg (25%) of 3-[3,5-dibromo-4-(3-ethylaminebenzyloxy)- phenyl]propionic acid. MS: m/z 457.2 (M^+-1).

Example 106**3-[3,5-Dibromo-4-(3-diethylaminebenzyloxy)phenyl]propionic acid**

Sodium boron hydride (17.6 mg, 0.47 mmol) was added in portions to a stirred solution of 3-[3,5-dibromo-4-(3-aminobenzyloxy)phenyl]propionic acid (40 mg, 0.090 mmol) in glacial acetic acid (5 mL) at room temperature. The reaction mixture was heated at 55 C for 2.5 hours. The reaction was cooled down to room temperature, quenched and purified on semi-preparative HPLC using the same conditions as described for Examples 27, 28 and 30. This gave 19 mg (42%) of 3-[3,5-dibromo-4-(3-diethylaminebenzyloxy)phenyl]propionic acid. MS: m/z 485.2 (M^+-1).

Example 107

N-[3,5-Dibromo-4-(3-bromobenzyloxy)phenyl]oxamic acid

(a) 3,5-Dibromo-4-hydroxynitrobenzene (40 mg, 0.13 mmol) was coupled with benzylbromide (0.26 mmol) using the procedure described in Method C. The crude material was used directly in the next step without additional purification on column. The crude material was dissolved in ethanol (3 mL) and treated with sodium dithionite (0.19 g, 1.1 mmol). After heating at 70°C over night, the reaction mixture was concentrated and dissolved in ethyl acetate. The organic phase was washed with an aqueous hydrochloric acid (0.01N), water and dried over $MgSO_4$. After concentration of the organic phase, the residue was treated with diethyl oxalate (0.5 mL). The reaction mixture was heated at 100 C for one hour and then at 180°C for 10 minutes. After cooling down to room temperature, the reaction mixture was purified on semi-preparative HPLC using the same conditions as described for Examples 27, 28 and 30. This gave 19 mg (26 %) of ethyl-N-[3,5-dibromo-4-(3-bromobenzyloxy)-phenyl]oxamate. MS: m/z 536.0 (M^+-1).

(b) Ethyl-N-[3,5-dibromo-4-(3-bromobenzyloxy)phenyl]oxamate (18 mg) was dissolved in methanol (1 mL) and treated with sodium hydroxide (1N, 6-8 equivalents). After stirring over night at room temperature, the reaction mixture was acidified to pH 5. The mixture was purified on semi-preparative HPLC using the same conditions as described for Examples 27, 28 and 30. This gave 7.8 mg (46 %) of N-[3,5-dibromo-4-(3-bromobenzyloxy)phenyl]oxamic acid. MS: m/z 508.0 (M^+-1).

Example 108

N-[3,5-Dibromo-4-(2-methylnaphthylloxy)phenyl]oxamic acid

3,5-Dibromo-4-hydroxynitrobenzene (40 mg, 0.13 mmol) was coupled with 2-(bromomethyl)naphthalene (0.26 mmol), reduced with sodium dithionite, coupled with diethyloxalate and hydrolyzed in a similar manner as described in Example 107. This gave 26 % N-[3,5-dibromo-4-(2-methylnaphthyloxy)phenyl]oxamic acid. MS: m/z 479.1 (M^+-1).

Example 109

3-[3-bromo-5-methoxy-4-(3-bromobenzyloxy)phenyl]propionic acid

Methyl 3-(3-bromo-4-hydroxy-5-methoxyphenyl) propionate (0.14 mmol) was coupled with benzylbromide (0.28 mmol) and hydrolyzed, using the procedure described for Examples 21-30 (Method C). This gave 15 % yield of 3-[3-bromo-5-methoxy-4-(3-bromobenzyloxy)phenyl]propionic acid. MS: m/z 444.1 (M^+-1).

Example 110

3-[3-bromo-5-methoxy-4-(2-methylnaphthyloxy)phenyl]propionic acid

Methyl-3-(3-bromo-4-hydroxy-5-methoxyphenyl) propionate (0.14 mmol) was coupled with 2-(bromomethyl)naphthalene (0.28 mmol) and hydrolyzed, using the procedure described for Examples 21-30 (Method C). This gave 18 % yield of 3-[3-bromo-5-methoxy-4-(2-methylnaphthyloxy)phenyl]propionic acid. MS: m/z 415.3 (M^+-1).

GENERAL PROCEDURE FOR THE PREPARATION OF EXAMPLES 111-119

To a solution of methyl 3-(3,5-dibromo-4-hydroxyphenyl) cinnamate (0.035 mmol) (prepared according to Schreiber, F. G; Stevenson, R., *J. Chem. Soc., Perkin Trans. I*, 1976, 14, 1514-18 and references cited therein) in dry acetonitrile (0.25 mL) was added potassium carbonate (0.070 mmol) and the resulting mixture was stirred at room temperature. After 10 minutes a solution of the halide (0.070 mmol) in acetonitrile (0.25 mL) was added. When the halide was a chloride, a catalytic amount of sodium iodide was added. The mixture was heated at 80 °C over night. After cooling down to room temperature, the reaction mixture was filtered through a silica SPE column (500 mg/3 mL), eluting with *n*-heptane/ethyl acetate 3:1 (3 mL). After concentration *in vacuo* the residue was dissolved in tetrahydrofuran (0.25 mL)


and lithium hydroxide (1 N in water, 0.25 mL) was added. The reaction mixture was stirred at room temperature. After 4 hours the mixture was neutralised on an SCX SPE column (500 mg/3 mL), using methanol or triethylamine (10% in methanol) as eluent.

The eluent was concentrated and the residue purified on a silica SPE column (500 mg/3 mL, heptane/ethyl acetate 9:1 (3 mL) followed by dichloromethane/methanol 9:1 (3 mL). The product containing fractions were collected and concentrated *in vacuo* to give the final product.










GENERAL PROCEDURE FOR THE PREPARATION OF EXAMPLES 120-121

A solution of methyl-3-(3-bromo-4-hydroxy-5-methoxyphenyl) cinnamate (0.14 mmol) (prepared according to Schreiber, F. G; Stevenson, R., *J. Chem. Soc., Perkin Trans. 1*, 1976, 14, 1514-18 and references cited therein) was coupled with the appropriate bromo compound (0.28 mmol benzylbromide or 2-(bromomethyl)- naphthalene) and hydrolyzed, using the procedure described for Examples 21-30 (Method C). This gave after purification on HPLC (using the same conditions as described for Examples 27, 28 and 30) the final products.

For a table of Examples 111-121 comprising yield and mass spectra , see Table below.




Example	R	R ₄	Yield (%)	MS ¹
111		Br	35	478.9
112		Br	34	424.9

113		Br	58	495.1
114		Br	28	412.0
115		Br	10	461.8
116		Br	53	497.2
117		Br	62	415.9
118		Br	47	488.8
119		Br	10	460.9
120		OMe	19	442.1
121		OMe	6	413.3

The compounds of the invention exhibit binding affinities to the ThRa receptor in the range of 100nM to 10000nM.

CLAIMS

1. Compounds with the general formula:



or pharmaceutically acceptable salts thereof, wherein:

R_1 is selected from: C_{5-10} aryl; C_{5-10} heteroaryl; C_{3-15} alkyl; C_{4-15} alkenyl; C_{3-15} alkynyl; and C_{3-10} cycloalkyl, said aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl optionally being substituted with 1 to 3 groups of R^a ;

R_2 is selected from: hydrogen; C_{1-4} alkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; and C_{1-4} alkoxy; or a bioisosteric equivalent;

R_1 may be linked through the available carbons to R_2 , thus forming a C_4-C_8 cycloalkyl, saturated or partially unsaturated, and optionally substituted with 1 to 3 groups of R^a ;

X is selected from: -O-; -S-; -SO-; SO_2 -; -Se-; -Te-; $-N(R^c)$ -; and -S-S-;

R_3 and R_4 are the same or different and are selected from: halogen; C_{1-4} alkyl; C_{3-7} cycloalkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; C_{1-4} alkoxy; $-CF_3$; $-OCF_3$; $-OCF_2H$; $-SCH_3$; $-SCF_3$; and $-COOH$, or a bioisosteric equivalent; with the proviso that when R_3 is C_{1-4} alkoxy; $-OCF_3$; $-OCF_2H$; $-SCH_3$; $-SCF_3$; or $-COOH$, then R_4 must be selected from: halogen; C_{1-4} alkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; C_{3-7} cycloalkyl; and $-CF_3$;

n is an integer of 0, 1, 2 or 3;

Y is selected from: of -CO-; -O-; -S-; -CH(R^b)-; and -N(R^c)-;

R₅ is independently selected from:

carboxylic acid (-CO₂H);

phosphonic acid (-PO(OH)₂);

phosphamic acid (-PO(OH)NH₂);

sulphonic acid (-SO₂OH);

α-keto carboxylic acid (COCO₂H);

hydroxamic acid (-CONHOH);

sulphonamide (-SO₂NHR');)

sulphonamide (-NHSO₂R'), with the proviso that then Y is CO and n is 0;

acylsulphonamide (-CONHSO₂R') or (-SO₂NHCOR');

or any other possible bioisosteric equivalent of all the groups above;

R^a represents a member selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; -SMe; -CN; -NO₂; and -N(C₀₋₄)₂, or a bioisosteric equivalent;

R^b represents a member selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; and -(CH₂)_mOH, where m is an integer between 0 and 4, or a bioisosteric equivalent;

R^c represents a member selected from: hydrogen; C₁₋₄ alkyl; C₃₋₄ alkenyl; and C₃₋₄ alkynyl or a bioisosteric equivalent;

included for the variables above are all the possible stereoisomers thereof; prodrug ester forms thereof; and radioactive forms thereof.

2. The compounds according to claim 1 wherein:
X is -O-; and
other variables are kept as in claim 1.
3. The compounds according to claim 1 wherein:

R_2 is selected from hydrogen and methyl;

X is -O-;

R_3 and R_4 are the same or different and are halogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, or $-CF_3$;

n is an integer of 0 or 1;

Y is $-CH(R^b)-$;

R_5 is a carboxylic acid ($-CO_2H$);

R^b is hydrogen; and

other variables kept as described in claim 1.

4. The compounds according to claim 1 wherein:

X is -O-;

Y is $-N(R^c)-$;

n is an integer of 0, 1 and 2;

R_3 is a carboxylic acid ($-CO_2H$) or an α -keto carboxylic acid ($COCO_2H$);

R^c is hydrogen; and

other variables kept as described in claim 1.

5. The compounds according to claim 1 wherein:

X is -O-;

Y is -O-;

n is an integer of 1 and 2;

R_3 is a carboxylic acid ($-CO_2H$); and

other variables kept as described in claim 1.

6. The compounds according to claim 1 wherein:

X is -O-;


Y is $-CO-$;

n is 0;














R_3 is a sulphonamide ($-NHSO_2R'$); and

other variables kept as described in claim 1.


7. The compounds according to claim 1 wherein:
 R_2 is selected from hydrogen and methyl;
 X is $-O-$;
 R_3 is $-COOH$ and R_4 is selected from: halogen; C_{1-4} alkyl; C_{2-4} alkenyl; C_{2-4} alkynyl;
 and $-CF_3$;
 R^b is hydrogen;
 R^c is hydrogen; and
 other variables kept as described in claim 1.
8. The compounds according to any one of claims 1 to 7 which are:




R	n
	1
	1
	1
	1
	1
	1
	1

	2
	2
	2
	2
	2
	2
	2
 Me	2
	2
	2
	2
	2
	1


59




1




2




2




2




2




2




2



2




2





2

N-[3,5-Dibromo-4-(3-bromobenzoyloxy)benzoyl]benzenesulphonamide;
N-[3,5-Dibromo-4-(3-bromobenzoyloxy)benzoyl]-3-nitrobenzenesulphonamide;
N-[3,5-Dibromo-4-(3-bromobenzoyloxy)benzoyl]-4-nitrobenzenesulphonamide;
 4-Amino-*N*-[3,5-dibromo-4-(3-bromobenzoyloxy)benzoyl]benzenesulphonamide;
N-[3,5-Dibromo-4-(3-bromobenzoyloxy)benzoyl]methanesulphonamide;
 and the compounds indicated in the table below:




R	R ₃ , R ₄	Y	n
	Br	CH ₂	0
	Br	CH ₂	0
	Br	CH ₂	1
	Br	CH ₂	1

Br CH₂ 1Br CH₂ 1

Cl CHOH 0



Cl CHOH 0




Cl CHOH 0




Cl C=O 0




Cl C=O 0



Cl C=O 0
























Cl C=O 0














Cl C=O 0


Me CH₂ 0


	Me	CH ₂	0
	Me	CH ₂	0
	Me	CH ₂	0
	Me	CH ₂	0
	Me	CH ₂	0
	Me	C=O	0
	Me	C=O	0
	Me	C=O	0
	Me	C=O	0
	Me	C=O	0
	<i>i</i> -Pr	C=O	0












3-[3,5-Dibromo-4-(3-ethoxybenzyloxy)phenyl]propionic acid;
 3-[3,5-Dibromo-4-(3-propyloxybenzyloxy)phenyl]propionic acid;
 3-[3,5-Dibromo-4-(3-butyloxybenzyloxy)phenyl]propionic acid;
 3-[3,5-Dibromo-4-(3-aminobenzyloxy)phenyl]propionic acid;
 3-[3,5-Dibromo-4-(3-ethylaminebenzyloxy)phenyl]propionic acid;
 3-[3,5-Dibromo-4-(3-diethylaminebenzyloxy)phenyl]propionic acid;
 N-[3,5-Dibromo-4-(3-bromobenzyloxy)phenyl]oxamic acid;
 N-[3,5-Dibromo-4-(2-methylnaphthyloxy)phenyl]oxamic acid;
 3-[3-bromo-5-methoxy-4-(3-bromobenzyloxy)phenyl]propionic acid;
 3-[3-bromo-5-methoxy-4-(2-methylnaphthyloxy)phenyl]propionic acid;
 and pharmaceutically acceptable salts thereof, and stereoisomers thereof.

9. A compound according to any one of claims 1 to 8, which have one or more asymmetric centers and can exist in the form of racemates, single and multiple enantiomers, as individual diastereomers, with all possible isomers, and mixtures thereof.
10. A compound according to any one of claims 1 to 9 for use in medical therapy.
11. A pharmaceutical composition comprising an effective amount of a compound according to any one of claims 1 to 9 or a pharmaceutically effective salt thereof, together with a pharmaceutically acceptable carrier.
12. A method for preventing, inhibiting or treating a disease which is dependent on the expression of a T₃ regulated gene or associated with metabolic dysfunction, which

comprises administering to a patient in need of treatment a therapeutically effective amount of a compound according to any one of claims 1 to 9.

13. The method according to claim 12 wherein the disease is selected from cardiac arrhythmias, thyrotoxicosis, subclinical hyperthyroidism and certain skin disorders.
14. The method according to claim 12 wherein the disease is a skin disorder or skin disease.
15. The method according to claim 14 wherein the skin disorder or skin disease is selected from: keloids, lichen planus, ichthyosis, acne, psoriasis, Dernier's disease, eczema, atopic dermatitis, chloracne, pityriasis, and hirsutism.
16. A method to treat certain skin disorders or diseases by the use of a compound according to any one of claims 1 to 9 in a combination with a retoid or a Vitamin D analog.
17. The use of a compound according to any one of claims 1 to 9 in the preparation of a medicament for the treatment of a disease or disorder which is dependent on the expression of a T₃ regulated gene.
18. The use according to claim 17 wherein the disease or disorder is cardiovascular disorder, thyrotoxicosis, subclinical hyperthyroidism and certain skin disorders.
19. The use according to claim 17 wherein the disease or disorder is selected from atrial fibrillation, ventricular tachycardia and ventricular fibrillation.
20. The use according to claim 17 wherein the disease or disorder is selected from thyrotoxicosis, subclinical hyperthyroidism and other related endocrine disorders, related to thyroid hormone.
21. The use of a compound according to any one of claims 1 to 9 in the preparation of a medicament for the treatment of anoxic tissue damage.

22. The use of a labeled compound, according to any one of claims 1 to 9, as a diagnostic agent.
23. The use of a compound in medical therapy according to the formula below:



wherein:

R_1 is selected from: of C_{5-10} aryl; C_{5-10} heteroaryl; C_{3-15} alkyl; C_{4-15} alkenyl; C_{3-15} alkynyl; and C_{3-10} cycloalkyl, said aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl optionally being substituted with 1 to 3 groups of R^a ;

R_2 is selected from: hydrogen; C_{1-4} alkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; and C_{1-4} alkoxy, or a bioisosteric equivalent;

R_1 may be linked through the available carbons to R_2 , thus forming a C_4-C_8 cycloalkyl, saturated or partially unsaturated, and optionally substituted with 1 to 3 groups of R^a ;

X is selected from: -O-; -S-; -SO-; SO_2 -; -Se-; -Te-; $-N(R^c)$ -; and -S-S-;

R_3 and R_4 are the same or different and are selected from: halogen; C_{1-4} alkyl; C_{3-7} cycloalkyl; C_{2-4} alkenyl; C_{2-4} alkynyl; C_{1-4} alkoxy; $-CF_3$; $-OCF_3$; $-OCF_2H$; $-SCH_3$; $-SCF_3$; $-COOH$ or a bioisosteric equivalent;

Y is selected from: $-CO-$; $-O-$; $-S-$; $-CH(R^b)-$; and $-N(R^c)-$;

R₃ is independently selected from:

carboxylic acid (-CO₂H);

phosphonic acid (-PO(OH)₂);

phosphamic acid (-PO(OH)NH₂);

sulphonic acid (-SO₂OH);

α-keto carboxylic acid (COCO₂H);

hydroxamic acid (-CONHOH);

sulphonamide (-SO₂NHR');)

sulphonamide (-NHSO₂R');

acylsulphonamide (-CONHSO₂R') or (-SO₂NHCOR'),

or any other possible bioisosteric equivalent of all the groups above;

R₆ and R₇ are the same or different and are selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; and -(CH₂)_mOH; where m is an integer between 1 and 4, or a bioisosteric equivalent;

R^a is selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; -SMe; -CN; -NO₂; and -N(C₀₋₄)₂, or a bioisosteric equivalent;

R^b represents a member selected from: hydrogen; halogen; -CF₃; C₁₋₄ alkyl; C₂₋₄ alkenyl; C₂₋₄ alkynyl; C₁₋₄ alkoxy; and -(CH₂)_mOH, where m is an integer between 0 and 4, or a bioisosteric equivalent;

R^c is selected from: hydrogen; C₁₋₄ alkyl; C₃₋₄ alkenyl; and C₃₋₄ alkynyl, or a bioisosteric equivalent;

or pharmaceutically acceptable salts thereof;

included for the variables above are all the possible stereoisomers thereof; prodrug ester forms thereof; and radioactive forms thereof.

24. The use of the compounds according to claim 23 wherein:

R_2 is selected from hydrogen and methyl;

X is -O-;








R_3 and R_4 are the same or different and are selected from: halogen; C_{1-4} alkoxy; C_{1-4} alkyl; and $-CF_3$;


R_5 is a carboxylic acid ($-CO_2H$);

R_6 and R_7 is hydrogen; and


other variables kept as described in claim 21.

25. The use of the compounds as defined in claims 23 to 24, said compounds being:







Br




Br




Br



Br



OMe



OMe

or pharmaceutically acceptable salts thereof and stereoisomers thereof.

26. The use of a compound according to any one of claims 23 to 25, wherein the compound have one or more assymetric centers and can exist in the form of racemates, single and multiple enantiomers, as individual diastereomers, with all possible isomers, and mixtures thereof.
27. The use of a compound according to any one of claims 23 to 26 in the preparation of a medicament for the treatment of a disease or disorder which is dependent on the expression of a T₃ regulated gene.

28. The use according to claim 27 wherein the disease or disorder is selected from atrial fibrillation, ventricular tachycardia, and ventricular fibrillation.
29. The use according to claim 27 wherein the disease or disorder is selected from thyrotoxicosis, subclinical hyperthyroidism, and other related endocrine disorders, related to thyroid hormone.
30. The use according to claim 27 wherein the disease or disorder is selected from cardiovascular disorder, thyrotoxicosis, subclinical hyperthyroidism and certain skin disorders.
31. The use of a compound according to any one of claims 23 to 26 in the preparation of a medicament for the treatment of anoxic tissue damage.
32. A pharmaceutical composition comprising an effective amount of a compound according to any one of claims 23 to 26 or a pharmaceutically effective salt thereof, together with a pharmaceutically acceptable carrier.
33. A method for preventing, inhibiting or treating a disease which is dependent on the expression of a T_3 regulated gene or associated with metabolic dysfunction, which comprises administering to a patient in need of treatment a therapeutically effective amount of a compound according to any one of claims 23 to 26.
34. The method according to claim 33 wherein the disease is selected from: cardiac arrhythmias, thyrotoxicosis, subclinical hyperthyroidism and certain skin disorders.
35. The method according to claim 33 wherein the disease or disorder is a skin disorder or skin disease.
36. The method according to claim 35 wherein the skin disorder or skin disease is keloids, lichen planus, ichthyosis, acne, psoriasis, Dernier's disease, eczema, atopic dermatitis, chloracne, pityriasis, eczema or hirsutism.

37. A method to treat certain skin disorders or diseases by the use of a compound according to any one of claims 23 to 26 in a combination with a retoid or a Vitamin D analog.
38. The use of a labeled compound, according to any one of claims 23 to 26, as a diagnostic agent.
39. A compound according to claim 10 for use in treating a disease or disorder that is dependent on the expression of a T_3 regulated gene or associated with metabolic disfunction.
40. A compound according to claim 39 wherein the disease is selected from cardiac arrhythmias, thyrotoxicosis, subclinical hyperthyrodism and skin disorders or skin diseases.
41. The compound according to claim 40 wherein the skin disorder or skin disease is selected from : keloids, lichen planus, ichtyosis, acne, psoriasis, Dernier's disease, eczema, atopic dermatitis, chloracne, pityriasis, and hirsutism.
42. A compound according to claim 39 wherein the disease or disorder is selected from atrial fibrillation, ventricular tachycardia and ventricular fibrillation.
43. A compound according to claim 39 wherein the disease or disorder is selected from thyrotoxicosis, and other related endocrine disorders, related to thyroid hormone.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number
WO 01/036365 A3

(51) International Patent Classification⁷: C07C 59/64,
59/72, 59/90, 205/37, 311/53, 255/54, C07D 213/30,
C07C 215/14, C07D 215/14, 333/56, 215/18, 261/08,
277/24, 309/06, A61P 9/06, 5/14, A61K 31/192, 31/277

Novum, S-141 57 Huddinge (SE). GORDON, Sandra
[UY/SE]; Karo Bio AB, Novum, S-141 57 Huddinge (SE).

(21) International Application Number: PCT/EP00/11554

(74) Agents: BANNERMAN, David, Gardner et al.; With-
ers & Rogers, Goldings House, 2 Hays Lane, London SE1
2HW (GB).

(22) International Filing Date:
16 November 2000 (16.11.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9927056.3 17 November 1999 (17.11.1999) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): KARO
BIO AB [SE/SE]; Novum, S-141 57 Huddinge (SE).

(72) Inventors; and

Published:
— with international search report

(75) Inventors/Applicants (*for US only*): MALM, Johan
[SE/SE]; Korsangarvagen 53, S-142 40 Skogas (SE). LIT-
TEN, Chris [NZ/SE]; Stubbestigen 6, S-147 52 Tumba
(SE). APELQVIST, Theresa [SE/SE]; Sagstuvagen 2E,
3tr, S-141 49 Huddinge (SE). HEDFORS, Asa [SE/SE];
Domarvagen 3nb, S-141 62 Huddinge (SE). BRANDT,
Peter [SE/SE]; Karo Bio AB, Novum, S-141 57 Huddinge
(SE). EDVINSSON, Karin [SE/SE]; Karo Bio AB,

(88) Date of publication of the international search report:
7 November 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/036365 A3

(54) Title: THYROID RECEPTOR ANTAGONISTS FOR THE TREATMENT OF CARDIAC AND METABOLIC DISORDERS

(57) Abstract: This invention relates to novel compounds which are thyroid receptor ligands, preferably antagonists, and to methods for using such compounds in the treatment of cardiac arrhythmias, thyrotoxicosis and subclinical hyperthyroidism.

INTERNATIONAL SEARCH REPORT

 International Application No.
 PCT/EP 00/11554

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 C07C59/64 C07C59/72 C07C59/90 C07C205/37 C07C311/53
 C07C255/54 C07D213/30 C07C215/14 C07D215/14 C07D333/56
 C07D215/18 C07D261/08 C07D277/24 C07D309/06 A61P9/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DIBBO, A., SLY, J.C.P., STEPHENSON, L., WALKER, T., WARBURTON, W.K., WHITING, K.D.E.: "The Synthesis of Thyroxine and Related Compounds. Part XVII. The Preparation of Some Additional Compounds Related to Thyroxine." THE JOURNAL OF THE CHEMICAL SOCIETY, no. 3, 1961, pages 2890-2902, XP002177006 page 2890, paragraph 1 page 2892, paragraph 4	1-3, 8, 10-13
Y	---	1-3, 8, 10-13
	-/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 September 2001

Date of mailing of the international search report

21/09/2001

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

DeLanghe, P

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 00/11554

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61P5/14 A61K31/192 A61K31/277

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BUU-HOI N P ET AL: "ACIDES ARYLACETHYDROXAMIQUES ANTI-INFLAMMATOIRES ET ANALGESIQUES" CHIMICA THERAPEUTICA, SOCIETE D'ETUDES DE CHIMIE THERAPEUTIQUE, FR, vol. 2, no. 1, 1967, pages 39-48, XP000573817 abstract table 2 page 47, column 2, paragraph 5	1-3, 10, 11
X	US 3 586 713 A (BUU-HOI NGUYEN PHUC ET AL) 22 June 1971 (1971-06-22) examples 1,7 column 1, line 63-71 claims 1,8,9	1-3, 10, 11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

14 September 2001

Date of mailing of the international search report

Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

DeLanghe, P

INTERNATIONAL SEARCH REPORT

In
 Application No
PCT/EP 00/11554

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 28137 A (PATCHETT ARTHUR A ;BERGER JOEL P (US); ADAMS ALAN D (US); FITCH KE) 7 August 1997 (1997-08-07) abstract page 108; example 41 page 91; example 30 page 93; example 31	1-3,10, 11
X	FR 5 837 M (BEECHAM GROUP LIMITED AND MONSANTO CHEMICALS LIMITED) 10 April 1968 (1968-04-10) page 1, column 1, paragraphs 1,3-5	1,2,4, 10,11
X	EP 0 257 828 A (SUMITOMO CHEMICAL CO) 2 March 1988 (1988-03-02) table 1, compounds 19-20	1-3,5
Y	US 3 789 070 A (ALDO G) 29 January 1974 (1974-01-29) the whole document, specifically the examples 4-8	1-3,8, 10-13
A	EP 0 512 352 A (HOFFMANN LA ROCHE) 11 November 1992 (1992-11-11) page 1-11	1,12,17
A	US 5 516 931 A (GIESE ROGER W ET AL) 14 May 1996 (1996-05-14) column 19-20 claims 4-6	23

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-5, 7, 9-24, 26-43 (partially searched); 6 (not searched at all)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims 1-2 may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims 1-2 is impossible. Consequently, the search has been restricted to the compounds of the formula in claim 1 (where R1, R2, R3, R4, n, Ra, Rb, Rc and Y are as indicated and) where R5 = COOH including the pharmaceutical salts, stereoisomers and radioactive forms thereof and where X = 0.

Present claims 1-4 and 9 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of the formula in claim 1 (where R1, R2, R3, R4, n, Ra, Rb, Rc and Y are as indicated and) where R5 = COOH including the pharmaceutical salts, stereoisomers and radioactive forms thereof and where X = 0.

The use claims 10, 12-22, 39-43 have been similarly limited.
The composition claim 11 has been similarly limited.

Present claims 23, 24, 26-31, 33-38 specifies the use in medical therapy of a series of compounds, defined by a generic formula given in claim 23.

This formula, which characterizes the subject matter of the claims, relates to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the subject matter claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the subject matter of the formula in claim 23 (where R1-R4, R6, R7, Ra, Rb, Rc are as indicated and) where R5 = COOH including the pharmaceutical salts, stereoisomers and radioactive forms thereof and where X = 0.

The composition claim 32 has been similarly limited.

The scope of claims 1-7, 23, 24, in as far as the expressions bioisosteric equivalent and prodrug ester forms is concerned, is so unclear (Article 6 PCT) that a meaningful International Search is

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

impossible with regard to this expression.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

 In: International Application No
 PCT/EP 00/11554

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3586713	A	22-06-1971	BE 689347 A	14-04-1967
			CH 475201 A	15-07-1969
			DE 1543784 A1	09-03-1972
			FR 6048 M	11-07-1968
			FR 1503394 A	08-02-1968
			GB 1166818 A	08-10-1969
			NL 6617354 A	12-06-1967
			NL 7017022 A	17-06-1971
			OA 2280 A	05-05-1970
			ES 334305 A1	01-02-1968
WO 9728137	A	07-08-1997	AU 708055 B2	29-07-1999
			AU 1856397 A	22-08-1997
			AU 1856997 A	22-08-1997
			EP 0882029 A1	09-12-1998
			WO 9728137 A1	07-08-1997
			WO 9728149 A1	07-08-1997
			US 6090836 A	18-07-2000
			ZA 9700824 A	30-10-1998
FR 5837	M	04-03-1968	BE 690489 A	30-05-1967
			CH 481066 A	15-11-1969
			DE 1543382 A1	07-08-1969
			DK 118395 B	17-08-1970
			GB 1174051 A	10-12-1969
			NL 6616594 A	31-05-1967
			US 3492349 A	27-01-1970
EP 0257828	A	02-03-1988	AU 592448 B2	11-01-1990
			AU 7677087 A	03-03-1988
			CA 1317977 A1	18-05-1993
			DE 3767084 D1	07-02-1991
			EP 0257828 A1	02-03-1988
			JP 8025944 B	13-03-1996
			JP 63239244 A	05-10-1988
			KR 9504029 B1	22-04-1995
			US 4870064 A	26-09-1989
US 3789070	A	29-01-1974	BE 768811 A1	03-11-1971
			CA 919686 A1	23-01-1973
			DE 2131677 A1	17-02-1972
			FR 2100828 A5	24-03-1972
			ZA 7104114 A	29-03-1972
EP 0512352	A	11-11-1992	US 5344843 A	06-09-1994
			AT 136018 T	15-04-1996
			AU 653398 B2	29-09-1994
			AU 1600392 A	12-11-1992
			BR 9201769 A	29-12-1992
			CA 2068076 A1	10-11-1992
			CZ 9201385 A3	17-02-1993
			DE 69209357 D1	02-05-1996
			EP 0512352 A2	11-11-1992
			FI 922095 A	10-11-1992
			HU 63602 A2	28-09-1993
			IE 921489 A1	18-11-1992
			JP 5279353 A	26-10-1993
			JP 7107060 B	15-11-1995

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/EP 00/11554

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0512352 A		MX 9202153 A1	01-11-1992
		NO 921840 A	10-11-1992
		NZ 242626 A	27-01-1995
		RO 109938 B1	28-07-1995
		RU 2059603 C1	10-05-1996
		ZA 9203279 A	27-01-1993
US 5516931 A	14-05-1996	US 5360819 A	01-11-1994
		US 4709016 A	24-11-1987
		US 5610020 A	11-03-1997
		US 5602273 A	11-02-1997
		US 5604104 A	18-02-1997
		CA 1246058 A1	06-12-1988
		DK 36483 A	02-08-1983
		EP 0085554 A1	10-08-1983
		ES 519424 D0	01-08-1984
		ES 8406736 A1	01-11-1984
		JP 58146540 A	01-09-1983
		US 5650270 A	22-07-1997
		US 4650750 A	17-03-1987

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.